

para posteriormente utilizarlo como elemento clave para buscar hongos que tengan este compuesto en una mayor concentración y ejerzan un mayor porcentaje de inhibición de *Attamyces* sp.

BIBLIOGRAFÍA

Askew, D. J., and Laing, M. D. The *in vitro* screening of 118 *Trichoderma* isolates for antagonism to *Rhizoctonia solani* and an evaluation of different environmental sites of *Trichoderma* as sources of aggressive strains. *Plant and Soil.* 1994. 159:277-281.

Bacci, M., Ribeiro, S. B., Casarotto, M. E., Pagnocca, F. C. Biopolymer - degrading bacteria from nest of the leaf - cutting ant *Atta sexdens rubripilosa*. *Brazilian Journal of Medical and Biological Research.* 1995. 28:79-82.

Bass, M., and Cherrett, J. M. Fungus garden structure in the leaf-cutting ant *Atta sexdens* (Formicidae, Attini). *Symbiosis.* 1996 (a). 21:9-24.

_____. Leaf cutting ants (Formicidae, Attini) prune their fungus to increase and direct its productivity functional. *Ecology.* 1996 (b). 10: 55-61.

_____. The role of leaf-cutting ant workers (Hymenoptera: Formicidae) in fungus garden maintenance. *Ecological Entomology.* 1994. 19:215-220.

Beale, R. E., and Pitt, D. The antifungal properties of *Minimedusa polyspora*. Mycol. Res. 1995. 99:337-342.

Bélager, R.R., Dufour, N., Caron, J., and Benhamou, N. Chronological events associated with the antagonistic properties of *Trichoderma harzianum* against *Botrytis cinerea*: indirect evidence for sequential role of antibiosis and parasitism. Bioc. Sci. and Tech. 1995. 5:41-53.

Benhamou, N., and Chet, I. Parasitism of Sclerotia of *Sclerotium rolfsii* by *Trichoderma harzianum*: ultrastructural and cytochemical aspects of the interaction. Biochem. and Cell Bio. 1996. 86:405-415.

Brener, A. F., and Ruggiero, A. Leaf - cutting ant (*Atta* and *Acromyrmex*) inhabiting Argentina: Patterns in species richness and geographical range sizea. Journal of Biogeography. 1994. 21:391-399.

Benyogoub, M., Jabaji-Here, S. H., Banville, G., Charest, P. M. *Stachybotrys elegans*: a destructive mycoparasite of *Rhizoctonia solani*. Mycol. Res. 1994. 98:493-505.

Bueno, O. C., Hebling, M. J. A., Silva, O. A., Matenhauer, A. M. C. Effect of sesame (*Sesamum indicum*) on nest development of *Atta sexdens rubropilosa* (Hymenoptera: Formicidae). J. Appl. Ent. 1995. 119:341-343.

Chapela, I. H., Rehner, S. A., Schultz, T. R., Mueller, U. G. Evolutionary history of the symbiosis between fungus-growing ants and their fungi. SCIENCE. 1994. 266:1691-1694.

Cherrett, J. M., Powell, R. J., and Stradling, D. J. The mutualism between Leaf-cutting ants and their fungus. insect - fungus Interactions. Royal Entomol. Soc. 1989. 14:93-120.

✓ Chet, I., and Inbar, J. Biological control of fungal pathogens. Appl. Biochem. and Biotechnol. 1994. 48:37-43.

Correa, A., Roquebert, M.F., and Bettucci, L. Trichorziainins activity on mycelial growth of *Sclerotium cepivorum* under laboratory conditions *in vitro*. Crytogramie, Mycol. 1996. 17:123-128.

Cotes, A. M. Study of common bean protection against damping - off by treatment of seeds with *Trichoderma koningii oudemans*. 1993, [Thesis] Faculte des Sciences Agronomiques de Gembloux, Unité de Pathologie Végétale. Belgique. 120p.

Da Silva, M. E., e Diehl-Fleig, E. Avaliação de diferentes linhagens de fungos entomopatogênicos para controle da Formiga *Atta sexdens piriventris* (Santschi, 1919) (Hymenoptera: Formicidae). An. Soc. Entomol. Brasil. 1988. 17:263-269.

Diehl-Fleig, E. Sucesso no estabelecimento de colônias de *Acromyrmex striatus* (Roger) (Hymenoptera: Formicidae). An. soc. entomol. Brasil. 1995. 24:625-630.

Diehl-Fleig, E., and Lucchese, de Paula, M. E. Nest foundation by *Acromyrmex striatus* (Hymenoptera, Formicidae). Biology and Evolution of social insects. 1992. p.51-54

Diehl-Fleig, E., da Silva, M. E., e Pacheco, M. M. R. Testes de patogenicidade dos fungos entomopatogênicos *Beauveria bassiana* e *Metarhizium anisopliae* em *Atta sexdens piriventris* (Santschi, 1919) em diferentes temperaturas. Ciência e cultura. 1988. 40:1103-1105.

Diehl-Fleig, E., da Silva, M. E., Specht, A., Bortolás, E. P., Emprego do fungo entomopatogênico *Beauveria bassiana* em: Iscas para o controle das formigas cortadeiras *Acromyrmex* spp. em floresta implantada de *Eucalyptus grandis*. Séptimo congresso florestal estadual nova prata, R.S. 1992. 2:1139-1150.

Diehl-Fleig, E., e da Silva, M. E. Development of *Beauveria bassiana* (Bals.) Vuill. And of *Metarhizium anisopliae* (Metsch) Sorok. In culture containing *Hovenia dulcis* extract. Acta Biologica Leopoldasia. 1992. 14:15-22.

Diehl-Fleig, E., e Lucchese, de Paula, M. E. Reações comportamentais de operárias de *Acromyrmex striatus* (Hymenoptera, Formicidae) na presença de fungo entomopatogênico. Revta bras. Ent. 1991. 35: 101-107.

Diehl-Fleig, E., e Valim-Labres, M. E. Fungi isolated from leaf-cutting ants *Atta sexdens piriventris* and *Acromyrmex heyeri* (Hymenoptera, Formicidae): Mucor spp. effects on *Bauveria bassiana* entomopathogen. Ciência e cultura. 1993. 45:142-143.

- Doi, S., and Mori, M. Antifungal properties produced by *Trichoderma* isolates from sawdust media of edible fungi against wood decay fungi. Mat. and Org 1994. 28: 143-151.
- Doi S, Yamada A, and Mori M. Antagonistic effects of *Trichoderma* ssp. Isolated from sawdust media of endible fungi against wood decay fungi. Mat. and Org. 1994. 28:131-141.
- Domsch, K. H., Gams, W., Anderson, T. H. Compendium of soil fungi. IHW. Verlag, Germany. 1993. p 368-377; 794-809.
- Elías, R. A., Arcos, O. S. Contribución al estudio del control biológico de *Fusarium oxysporum*, *Rhizoctonia solani* y *Pythium* sp. Mediante diferentes especies de *Trichoderma*. Santa Fe de Bogotá, 1984, 72p. [Tesis], Departamento de Biología, Facultad de Agronomía, Universidad Nacional de Colombia.
- Enciclopedia de la ciencia y de la técnica. eds. Danae, S.A- Barcelona. 1977. 5: 2332.
- Fisher, P.J., Stradling, D.J., and D. N. Pegler. *Leucoagaricus basidiomata* from a live nest of the leaf - cutting ant *Atta cephalotes*. Mycol. Res. 1994. 98: 884-888.
- Fisher, P.J., Stradling, D.J., Sutton, B.C., and Petrini, L.E. Microfungi in the fungus gardens of the leaf-cutting ant *Atta cephalotes*: a primary study. Mycol. Res. 1996. 100:541-546.

García, C. A., Lapointe, S. L., Serrano, M. S. Métodos culturales para el control de la hormiga arriera *Acromyrmex landolti* (Forel) en los Llanos Orientales. Sociedad Colombiana de Entomología. Resumenes XVI Congreso. Medellín. 1989. p37.

Goldman, G. H., Hayes, C., Harman, G. E. Molecular and cellular biology of biocontrol by *Trichoderma* spp. Forum, TIBTECH. 1994. 12:478-482.

Hasegawa, E. Caste specialization in food storage in the dimorphic ant *Colobopsis nipponicus* (Wheeler). Ins. Soc. 1993. 40:261-271.

Hebling, M.J., Maroti, P.S., Bueno, O.C., Da Silva, O.A., Pagnocca, F.C. Toxic effects of leaves of *Ricinus communis* (Euphorbiaceae) to laboratory nests of *Atta sexdens rubropilosa* (Hymenoptera: Formicidae). Bull. Entomol. Res. 1996. 86:253-256.

Heller, W. E., and Theiler-Hedrich, R. Antagonism of *Chaetomium globosum*, *Gliocladium virens* and *Trichoderma viride* to four soil-borne Phytophthora species. J. Phytopathol. 1994. 141:390-394.

Hölldobler, B., and Wilson, E. O. The ants. Belknap Press of Harvard University Press, Cambridge. 1990. 615P.

Horvath, E.M., Bürgel, J.L., Messner, K. The production of soluble antifungal metabolites by the biocontrol fungus *Trichoderma harzianum* in connection with the formation of conidiopores. Mat. Org. 1995. 29:1-14.

Inbar, J., Mendez, A., and Chet, I. Hyphal interaction between *Trichoderma harzianum* and *Sclerotinia sclerotiorum* and its role in biological control. *Soil Biol. Biochem.* 1996. 28:757-763.

Jeffries, P., and Young, T.W.K. *Interfungal parasitic relationships.* eds. CAB INTERNATIONAL, Inglaterra. 1994. 296 p.

Kay, S. J., and Stewart, A. Evaluation of fungal antagonists for control of onion white rot in soil box trials. *Plant Pathology.* 1994. 43:371-377.

Lorito, M., Woo, S. L., D'Ambrosio, M., Harman, G. E., Hayes, C. K., Kubicek, C. P., and Scala, F. Synergistic interaction between cell wall degrading enzymes and membrane affecting compounds. *Molecular Plant-Microbe Interactions.* 1996. 9:206-213.

Luciano, M. E., Diehl-Fleig, E., e da Silva, M. E. Organismos associados a uma colônia de *Acromyrmex heyeri* (Hymenoptera: Formicidae) mantida em laboratório. *Acta Biologica Leopoldensia.* 1995. 17:47-56.

Machado, V., Diehl-Fleig, E., da Silva, M. E., Lucchese, de Paula, M. E. Reações observadas em colônias de algumas espécies de *Acromyrmex* (Hymenoptera- Formicidae) quando inoculadas com fungos entomopatogênicos. *Ciência e cultura.* 1988. 40:1106-1108.

McAllister, C.B., García-Romero, I., Godeas, A., and Ocampo, J.A. *In vitro* interactions between *Trichoderma koningii*, *Fusarium solani* and *Glomus mosseae*. *Soil Biol. Bioche.* 1994. 26:1369-1374.

Osando, J.M., and Waudo, S.W. Interaction between *Trichoderma* species and *Armillaria* root rot fungus of tea in Kenya. Interna. J. Pest Manag. 1994. 40:69-74.

Ousley, M. A., Lynch, J. M., Whipps, J. M. Potential of *Trichoderma* spp. as consistent plant growth stimulators. Bio. Fertil. Soils. 1994. 17:85-90.

Pagnocca, F.C., Da Silva, O.A., Hebling-Beraldo, M.J., Bueno, O.C. Toxicity of Sesame extracts to the symbiotic fungus of leaf-cutting ants. Bull. Entomol. Res. 1990. 80:349-352.

Pagnocca, F.C., Ribeiro, S.B., Torkomian, V.L., Hebling, M.J., Bueno, O.C., Da Silva, O.A., Fernandes, J.B., Vieira, P.C., Da Silva, M.F., Ferreira, A.G., Toxicity of lignans to symbiotic fungus of leaf- cutting ants. J. Chem. Ecol. 1996. 22:1325-1329.

Powell, R.J., and D.J. Stradling. factors influencing the growth of *Attamyces bromatificus*, a symbiont of *Attine* ants. Trans. Mycol. Soc. 1986. 87:205-213.

Pratt, S. C., Carlin, N. F., and Calabi, P. Division of labor in *Ponera pennsylvannica* (Formicidae: Ponerinae). Ins. Soc. 1994. 41:43-61.

Reaves, J. L., and Crawford, H. R. *In vitro* antagonism by *Ulocladium botrytis* of *Phellinus weiri*, *Heterobasidion annosum*, and *Armillaria ostoyae*. Eur. J. For. Path. 1994. 24:364-375.

Rincón, G., Leguizamón, J. E., Arbeláez, T. G. Control biológico de *Rhizoctonia solani* con *Trichoderma* spp. en semilleros de café. Revista CENICAFE. 1992. 43:73-83.

Samuels, G. J. *Trichoderma: a review of biology and systematics of the genus*. Mycol. Res. 1996. 100:923-935.

Score, A. J., and Palfreyman, J. W. Biological control of the dry rot fungus *Serpula lacrymans* by *Trichoderma* species: The effects of complex and synthetic media on interactions and hyphal extension rates. International Blodeterioration & Biodegradation. 1994. 33:115-128.

Seifert, K. A., Samson, R. A., Chapela, I. H. *Escovopsis aspergilloides*, a rediscovered hyphomycete from leaf-cutting ant nest. Mycologia. 1995. 87:407-413.

Specht, A., Diehl-Fleig, E., e da Silva, M. E. Atratividade de iscas de *Beauveria bassiana* (Bals.) Vuill. A formigas do gênero *Acromyrmex* (Hymenoptera; Formicidae). An. Soc. Entomol. Brasil. 1994. 23:99-104.

Statsoft. Statistic/Mac. Release 3, eds., Inc. United States of America, 1992.

Steinmetz, J., and Schönbeck, F. Conifer bark as growth medium and carrier for *Trichoderma* and *Gliocladium roseum* to control *Pythium ultimum* on pea. J. Plant . Dis. and Protec. 1994. 101:200-211.

Stradling, D. J. An introduction to the fungus-growing ants, Atti. P 15-18. En: Ant- Plant Interactions. Ed. Hulex, C. R and Cutler, D. F. Oxford Science

Publications. 1991.

Stradling, D. J., and Van Breda, J. M. Mechanisms affecting load size determination in *Atta cephalotes* L. (Hymenoptera, Formicidae). Ins. Soc. 1994. 41:423-434.

Vasconcelos, H. L., and Cherrtt, J. M. Changes in leaf - cutting ant populations (Formicidae: Attini) after the clearing of mature forest in Brazilian Amazonia. Studies on Neotropical Fauna and Environment. 1995. 30:107-113.

Weber, N. A. Gardening ants the Attines. eds . The American philosophical, Society. United States. 1972. 146 p.

Wetterer, J. K. Forager polymorphism, size - matching, and load delivery in the leaf - cutting ant, *Atta cephalotes*. Ecological Entomology. 1994 (b). 19:57-64.

_____. Ontogenetic changes in forager polymorphism and foraging ecology in the leaf - cutting ant *Atta cephalotes*. Oecologia. 1994 (a). 98:235-238.

Wilson, E. O. The insect societies. Belknap Press of Harvard University Press, Cambridge. 1971. 548P.

Zhang, J., Howell, C. R., and Starr, J. L. Supression of *Fusarium* colonization of cotton roots and *Fusarium* witt by seed tretments with

Gliocladium virens and *Bacillus subtilis*. Biocontrol Science and Technology. 1996. 6:175-187.